

## Erratum to: Statistical power to detect multiple paternity in populations of highly fertile species: how many females and how many offspring should be sampled?

David Veliz<sup>1</sup> · Pierre Duchesne<sup>2</sup> · Noemi Rojas-Hernandez<sup>1</sup> · Luis M. Pardo<sup>3,4</sup>

Published online: 19 December 2016  
© Springer-Verlag Berlin Heidelberg 2016

**Erratum to: Behavioral Ecology and Sociobiology**  
DOI 10.1007/s00265-016-2240-x

There was an error in the equation in the original version of this article. The correct equation is:

$$P(\text{case 2}) = C_{NL}^{NM2} / C_{NL}^{(NM1+NM2)}$$

Summing over the two independent cases, we finally obtain:

$$P_{\text{SPiMB}} = (C_{NL}^{NM1} + C_{NL}^{NM2}) / C_{NL}^{(NM1+NM2)}$$

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00265-016-2240-x>.

---

✉ David Veliz  
dveliz@uchile.cl

<sup>1</sup> Departamento de Ciencias Ecológicas, Instituto de Ecología y Biodiversidad (IEB), Núcleo Milenio en Ecología y Manejo Sustentable (ESMOI), Universidad de Chile, Santiago, Chile

<sup>2</sup> Département de Biologie, Université Laval, Québec, Canada

<sup>3</sup> Instituto de Ciencias Marinas y Limnológicas, Universidad Austral de Chile, Valdivia, Chile

<sup>4</sup> Centro de Investigación de Dinámica de Ecosistemas Marinos de Altas Latitudes (IDEAL), Valdivia, Chile